

Version No.			
3	0	9	1

ROLL NUMBER					



- 0  0  0  0  
 1  1  1  1  
 2  2  2  2  
 3  3  3  3  
 4  4  4  4  
 5  5  5  5  
 6  6  6  6  
 7  7  7  7  
 8  8  8  8  
 9  9  9  9

- 0  0  0  0  0  0  
 1  1  1  1  1  1  
 2  2  2  2  2  2  
 3  3  3  3  3  3  
 4  4  4  4  4  4  
 5  5  5  5  5  5  
 6  6  6  6  6  6  
 7  7  7  7  7  7  
 8  8  8  8  8  8  
 9  9  9  9  9  9

Answer Sheet No. \_\_\_\_\_

Sign. of Candidate \_\_\_\_\_

Sign. of Invigilator \_\_\_\_\_

Section – A is compulsory. All parts of this section are to be answered on this page and handed over to the Centre Superintendent. Deleting/overwriting is not allowed. Do not use lead pencil.

**CHEMISTRY HSSC-I**  
**SECTION – A (Marks 17)**  
**Time allowed: 25 Minutes**

حصہ اول لازمی ہے۔ اس کے جوابات اسی صفحہ پر دسے کر ناظم مرکز کے حوالے کریں۔ کاٹ کر دوبارہ لکھنے کی اجازت نہیں ہے۔ ایڈمنسٹریٹو کا استعمال ممنوع ہے۔

ہر سوال کے سامنے دیے گئے درست دائرہ کو پر کریں۔

Fill the relevant bubble against each question:

- The number of Hydrogen molecules required to form 36g of water ( $H_2O$ ):  
 $2H_2 + O_2 \longrightarrow 2H_2O$   
  $2.408 \times 10^{24}$       $24.08 \times 10^{24}$       $1.204 \times 10^{24}$       $6.02 \times 10^{23}$
- The values of Balmer series wave number lines lie in:  
 U.V     I.R     Visible     Microwaves
- Which of the following sets of quantum numbers is **incorrect** for an electron?  
  $n=2, l=0, m=0$       $n=3, l=1, m=1$       $n=3, l=2, m=1$       $n=4, l=1, m=2$
- Hybridization of Beryllium in  $BeCl_2$  will be:  
  $SP^3$       $SP^2$       $SP$       $dSP^2$
- Which one of the following gas molecules will only show translational motion?  
  $H_2$       $NH_3$       $He$       $CO_2$
- According to Kinetic Molecular Theory:  
  $v \propto \sqrt{m}$       $v \propto m$       $v \propto \sqrt{\frac{1}{m}}$       $v \propto \frac{1}{m}$
- Osmotic pressure ( $\pi$ ) does **NOT** depend on:  
 Molarity     Universal gas constant     Temperature     Radius
- Equilibrium constants  $K_c = K_p$ , when  $\Delta n$  is:  
 1     -1     Zero     -2
- $K_a$  is acid association constant, large  $PK_a$  value for an acid indicates that acid is:  
 Strong     Weak     Moderate     Water soluble

10. Which of the following is an acidic salt?   $\text{NaHCO}_3$    $\text{NH}_4\text{Cl}$    $\text{CH}_3\text{COONa}$    $\text{K}_2\text{SO}_4$
- 
11. Rate of reaction independent to concentration of reactants in:  Zero order reactions  First order reactions  2<sup>nd</sup> order reactions  3<sup>rd</sup> order reactions
- 
12. Which of the following gases will be most non-ideal at  $-10^\circ\text{C}$ ?   $\text{CO}_2$    $\text{H}_2$    $\text{N}_2$    $\text{NH}_3$
- 
13.  $\Delta H = \Delta E + P\Delta V$  is formula of:  Enthalpy  Work  Surrounding  Internal energy
- 
14. Which of the following relationship is incorrect?   $\Delta H_v > \Delta H_f$    $\Delta H_f > \Delta H_v$    $\Delta H_s > \Delta H_f$    $\Delta H_s > \Delta H_v$
- 
15. Oxidation state of 'O' in  $\text{KO}_2$  is:  -1  -2  -4   $-\frac{1}{2}$
- 
16. Which of the following has strongest intermolecular forces of attraction?   $\text{H}_{2(g)}$    $\text{Cl}_{2(g)}$    $\text{I}_{2(s)}$    $\text{CH}_{4(g)}$
- 
17. Lattice energy may also be called:  Affinity energy  Crystal energy  Bond energy  Ionization energy

**SUPPLEMENTARY TABLE**

Atomic No	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Symbol	H	He	Li	Be	B	C	N	O	F	Ne	Na	Mg	Al	Si	P	S	Cl	Ar	K	Ca
Mass No	1	4	7	9	11	12	14	16	19	20	23	24	27	28	31	32	35.5	40	39	40

—1HA-I 2209-3091 (L)—

ROLL NUMBER					



# CHEMISTRY HSSC-I

24

Time allowed: 2:35 Hours

Total Marks Sections B and C: 68

NOTE: Answer any FOURTEEN parts from Section 'B' and attempts any TWO questions from Section 'C' on the separately provided answer book. Write your answers neatly and legibly.

## SECTION - B (Marks 42)

Q. 2 Answer any FOURTEEN parts from the following. All parts carry equal marks. (14 x 3 = 42)

- (i) The liquid  $CHBr_3$  has a density of  $2.89 \text{ g/cm}^3$ . What volume of this liquid should be measured to contain a total of  $4.8 \times 10^{24}$  molecules of  $CHBr_3$  (M.Wt, C=12, H=1, Br=80)
- (ii) Point out the three defects of Bohr's model.
- (iii) How dipole moment help to determine the polarity of molecules? Apply this concept to determine the nature of  $CO_2$  and Cis-1, 2-dichloro ethene.
- (iv) Predict and draw the shape and bond angles of following molecules on the basis of VSEPR theory:  
(i)  $H_2S$  (ii)  $SnCl_2$  (iii)  $PCl_3$
- (v) Briefly explain azimuthal quantum number. How it helps to determine number of  $e^-$  in a subshell?
- (vi) Prove that absolute temperature of a gas is the measure of average kinetic energy of its molecules.  $K \cdot E \propto T$
- (vii) How the molar mass and density of a gas can be determined with the help of general gas equation?
- (viii) Why butane is gas at room temperature while hexane is liquid?
- (ix) Differentiate between Isomorphism and polymorphism with suitable examples.
- (x) Describe electron sea theory. How it explains the properties of metals?
- (xi)  $N_2 + 3H_2 \rightleftharpoons 2NH_3$   $K_c = 6 \times 10^{-1}$  at  $500^\circ C$  Predict the direction in which the system will shift to attain equilibrium when concentrations of  $H_2$ ,  $N_2$  and  $NH_3$  are  $1.0 \times 10^{-2} M$ ,  $1.0 \times 10^{-3} M$ ,  $1.0 \times 10^{-3} M$  respectively.
- (xii) Calculate the  $pH$  of a buffer when molar concentrations of  $NH_4OH$  and  $NH_4Cl$  are  $1.0M$  and  $0.1M$  respectively.  $PK_b$  of  $NH_4OH$  is  $4.75$ .
- (xiii) Explain with chemical equation why aqueous solution of:  
(i)  $NH_4Cl$  is acidic (ii)  $K_2CO_3$  is basic (iii)  $Na_2SO_4$  is neutral
- (xiv) Consider the following reaction  $2H_2 + 2NO \longrightarrow N_2 + 2H_2O$   $R = K[H_2][NO]^2$  if this reaction occurs in two steps then write its mechanism and predict the reaction intermediate.
- (xv) What is diffusion? Also state Graham's law of effusion and diffusion with mathematical expression.
- (xvi) Calculate the molality of 30%  $\frac{w}{w}$  solution of fructose ( $C_6H_{12}O_6$ ).
- (xvii) Define system, surroundings and boundary with a suitable example.
- (xviii) Predict the feasibility of the following reaction  $Sn + Mg^{2+} \longrightarrow Sn^{2+} + Mg$   $E^\circ_{Sn} = -0.14V$ ,  $E^\circ_{Mg} = -2.38V$
- (xix) Distillation under reduced pressure is often used for purification of sensitive liquids. Describe the process giving reason.
- (xx) Apply  $n+l$  rule and pick the orbital with the lower energy from each of the given pairs:  
(i)  $3d, 4s$  (ii)  $2p, 3s$  (iii)  $6p, 4s$

## SECTION - C (Marks 26)

Note: Attempt any TWO questions. All questions carry equal marks. (2 x 13 = 26)

- Q. 3 a. Consider the following reaction  $CH_4 + H_2O \longrightarrow CO + 3H_2$  (06)
- (i) What is the amount of  $CO$  produced if 30g of  $CH_4$  and 50g of  $H_2O$  is used
- (ii) In an experiment 22g of  $CO$  were produced, what is percentage yield?
- b. Describe construction of lead storage battery and reactions taking place during charging and discharging. (07)
- Q. 4 a. What is orbital hybridization? Explain the structure of  $HC \equiv CH$ ,  $BF_3$  and  $CH_4$  on the basis of hybridization. (06)
- b. State Le-Chatelier's principle. Briefly discuss the effect of increase in pressure, increase in concentration of  $SO_2$ , increase in temperature and increase in  $NO_2$  catalyst when following reaction is at equilibrium.  $2SO_2 + O_2 \xrightleftharpoons{NO_2(g)} 2SO_3(g)$   $\Delta H = -256 \text{ kJ/mol}$  (07)
- Q. 5 a. Draw complete Born Haber cycle for the formation of  $MgO$  from the following data. (06)
- $\Delta H_f^\circ$  of  $MgO = -602 \text{ kJ/mol}$ ,  $\Delta H_s^\circ$  of  $Mg = 150 \text{ kJ/mol}$ ,  $\Delta H_{I.E}^\circ$  of  $Mg^{2+} = 2180 \text{ kJ/mol}$ ,  
 $\Delta H_{at}^\circ$  of  $O_2 = 24 \text{ kJ/mol}$ ,  $\Delta H_{E.A}^\circ$  of  $O^{-1} = -141 \text{ kJ/mol}$ ,  $\Delta H_{E.A}^\circ$  of  $O^{-2} = 878 \text{ kJ/mol}$
- b. Why addition of solute increases the boiling point of solution? Explain quantitative aspects of elevation of boiling point and prove that  $\Delta T_b$  is inversely proportional to molar mass of solute. (07)

### SUPPLEMENTARY TABLE

Atomic No	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Symbol	H	He	Li	Be	B	C	N	O	F	Ne	Na	Mg	Al	Si	P	S	Cl	Ar	K	Ca
Mass No.	1	4	7	9	11	12	14	16	19	20	23	24	27	28	31	32	35.5	40	39	40

— 1HA-I 2209 (L) —